

Dispersion Quark Model of Heavy Meson Decays

Dmitri Melikhov *

Nuclear Physics Institute, Moscow State University, Moscow, Russia

We discuss applications of the dispersion quark model to exclusive semileptonic decays of heavy mesons. The transition form factors in this relativistic formulation of the quark model are given by double spectral representations through the wave functions of the initial and final mesons both in the scattering and the decay regions. An important feature of the model is that the form factors have the correct behavior in accordance with QCD in the case of meson decays induced by heavy parent quark transition both to heavy and light daughter quarks. We demonstrate that the dispersion quark model gives results compatible with the lattice calculations at large q^2 and thus provides reliable form factors in the whole decay region.

Theoretical description of hadronic amplitudes of the quark currents is one of the key problems of particle physics as such amplitudes provide a bridge between QCD formulated in the language of quarks and gluons and observable phenomena which deal with hadrons. In particular, the knowledge of such amplitudes is necessary for extraction of the parameters of the quark-mixing matrix in the Standard Model from the experiments on weak heavy hadron decays. The difficulty in such calculations lie in the fact that hadron formation occurs at large distances where perturbative QCD methods are not applicable and nonperturbative consideration is necessary.

Various theoretical frameworks more or less directly related to QCD have been applied to the description of meson transition form factors: among them lattice QCD [1], QCD sum rules [2], and constituent quark models [3].

Lattice QCD simulations is the most direct QCD based nonperturbative approach and thus should provide most reliable results. Although it is not possible to place b quark directly on the lattice, a constrained extrapolation in heavy quark mass [1] allows one to reliably determine the form factors for B meson decay. A more serious problem is that lattice calculations do not provide the form factors in the whole accessible kinematical region: the daughter light quark produced in the b decay cannot be allowed to move fast enough on the lattice and one finds himself constrained in the region of not very large recoils. For obtaining form factors in the whole kinematical decay region one can use the extrapolation procedures based on some parametrizations of the form factors. For instance, in [1] a simple lattice-constrained parametrization based on the constituent quark picture [4] and pole dominance is proposed. Anyway, a reliable knowledge of form factors in some region is already a substantial step forward which provides firm constraints on results of other approaches.

QCD sum rules give in a sense a complementary information on the form factors as they allow one to determine

the form factors at not very large momentum transfers [2]. However in practice various versions of QCD sum rules give rather uncertain predictions strongly dependent on the technical subtleties of the particular version. A recent analysis [5] disregards the three-point sum rules in favour of the light-cone sum rules. On the other hand, the light-cone sum rules involve more phenomenological inputs and the results turn out to be sensitive to a particular distribution amplitude of the light meson used in the evaluation of the sum rule and a model of subtracting the continuum (cf. [6] and [7]).

Constituent quark models have proved to be a fruitful phenomenological method for the description of heavy meson transitions (a detailed review of the quark model is given in the talk by A. Le Yaouanc [3]). An attractive feature of the approaches based on the concept of constituent quarks is that these approaches suggest a physical picture of the process. The physical picture of the constituent quark model is based on the following phenomena responsible for the soft physics: the chiral symmetry breaking in the soft region which yields appearance of constituent quarks and gluons; then, a strong peaking of the soft (nonperturbative) wave functions of hadrons with a width of order of the confinement scale; and, finally, the relevance of the contribution of only Fock state components with a minimal number of the constituents (i.e. $q\bar{q}$ component in mesons and qqq component in baryons). The latter assumption is justified by the successes of the constituent quark model of hadron spectrum. We would like to illustrate that basically these features of the soft physics yield quite reasonable quantitative results for the meson transition form factors.

An adequate framework which implements the above-mentioned features of the nonperturbative physics is the dispersion formulation of the quark model of meson decays [8] which is based on taking into account only two-particle $\bar{q}q$ intermediate states in the amplitudes of the $\bar{q}_s q \rightarrow \bar{q}_s q'$ transition through a bilinear quark current $\bar{q}' O q$.

The transition form factors of the meson M_1 to the meson M_2 both in the scattering and the decay region are given by the double relativistic spectral representations through the soft wave functions of the initial and final

*Talk given at the IVth International Workshop on Progress in Heavy Quark Physics, Rostock, September 20-22, 1997

mesons, $G_1(s_1)$ and $G_2(s_2)$, respectively

$$f(q^2) = \int \frac{ds_1}{s_1 - M_1^2} \frac{ds_2}{s_2 - M_2^2} G_1(s_1) G_2(s_2) \times \left[\tilde{f}_D(s_1, s_2, q^2) + \tilde{f}_{\text{sub}}(s_1, M_1^2, s_2, M_2^2, q^2) \right], \quad (1)$$

where s_1 (s_2) is the invariant mass of the initial (final) $\bar{q}q$ pair. The double spectral density of the representation (1) consists of the two parts: the unsubtracted part $\tilde{f}_D$ and the subtraction part $\tilde{f}_{\text{sub}}$.

The unsubtracted part $\tilde{f}_D(s_1, s_2, q^2)$ is calculated unambiguously from the double cut triangle Feynman graph with a relevant spinorial structure of the vertices corresponding to the initial and final mesons and to the quark transition current [8].

The subtraction part $\tilde{f}_{\text{sub}}(s_1, M_1^2, s_2, M_2^2, q^2)$ contains the factors $s_1 - M_1^2$ or $s_2 - M_2^2$ and accounts for an ambiguity connected with possible subtractions in the dispersion representations inherent to any dispersion approach. To decide on the necessity and the structure of the subtraction terms we need some additional arguments.

Notice that once we are working within an approach not directly deduced from QCD it is important to keep essential features of the underlying fundamental theory in the model. Thus matching the results obtained within the quark model to rigorous QCD results might be helpful for bringing more realistic features to the model.

QCD gives rigorous predictions for the structure of the form factors in the case of the meson transition induced by the heavy quark transition. And it turns out possible to use matching the form factors of the dispersion quark model to form factors in QCD in the heavy quark limit for determining the subtraction terms in the spectral representations (1).

Namely, we consider the case of a pseudoscalar heavy meson decay into final pseudoscalar and vector mesons induced by the heavy-to-heavy and heavy-to-light quark transitions through vector, axial-vector and tensor currents. We perform the expansion of the quark model form factors in inverse powers of the heavy quark mass m_Q and match this expansion to the heavy quark expansion in QCD in leading and next-to-leading orders [9]. This matching provides constraints on the structure of the subtraction terms. Additional constraints allowing us to fix the subtraction terms are obtained by requiring the form factors of vector, axial-vector and tensor currents to obey the general relations found in [10] in the case of heavy-to-light transitions.

It is important that the only property of the soft wave function of a heavy meson necessary for obtaining the correct $1/m_Q$ expansion of the transition form factors is a strong peaking of this soft wave function in the relative momenta of the constituent quarks in the region of order Λ_{QCD} . No other constraints on the soft wave function are required.

Finally, we arrive at the double spectral representations of the form factors with fixed subtraction terms.

To obtain numerical predictions for the form factors we must specify the parameters of the quark model, i.e. constituent quark masses and the meson wave functions. To this end we take as an example the parameters of the ISGW2 model [12]. The form factors of the semileptonic decay of B and D mesons calculated within dispersion quark model adopting the ISGW2 quark masses and wave functions provide reasonable description of all experimental data [11]. Fig.1 illustrates the form factor $f_+^{B \rightarrow \pi}$ vs the lattice results, and Fig. 2 presents form factors for the transition $B \rightarrow K^*$.




FIG. 1. The form factor $f_+^{B \rightarrow \pi}$. QM result-solid, LCSR [2] - dashed, lattice points from [1].




FIG. 2. $B \rightarrow K^*$ form factors. QM results: solid- T_2 , long-dashed- A_1 , dashed- T_1 , dash-dotted- A_0 , dotted- V . Lattice data from [1].

One can observe a reasonable agreement with the lattice results. For the value $T_2(0) = T_1(0)$ relevant for the

$B \rightarrow K^* \gamma$ transition we find

$$T_2(0) = 0.14.$$

which agrees well with a recent light-cone sum rule estimate [7].

It should be understood that the reported form factors are obtained with a very simple exponential ansatz for soft wave functions, which can be expected to describe only the effects of the confinement scale. Figs. 1 and 2 show that even this rather crude approximation works well and suggest that the bulk of the form factor behavior can be explained by the chiral symmetry breaking in the soft region and a strong peaking of the meson wave functions with a width of order of the confinement scale [4].

For obtaining more reliable predictions for the form factors one needs realistic meson wave functions and constituent quark masses relevant for the meson decays. These can be found by applying a combined fit based on the representations (1) to all available lattice points on meson transition form factors at large q^2 considering the wave functions and constituent quark masses as variational parameters. In this case the spectral representations of the dispersion quark model may be treated as some generalized parametrizations of the transition form factors based on the constituent quark picture and obeying the rigorous QCD results in the limit of heavy-to-heavy and heavy-to-light meson transitions.

The work reported here was performed in part in collaboration with N. Nikitin and S. Simula and supported by the RFBR under grant 96-02-18121a. I take pleasure in thanking D. Becirevic, M. Beyer, V. Braun and A. Le Yaouanc for helpful discussions and the Organizers for the invitation to this pleasant and stimulating meeting.

-
- [1] C. T. Sachrajda, these proceedings; J. M. Flynn and C. T. Sachrajda, e-print [hep-lat/9710057](#).
 - [2] V. M. Braun, these proceedings and refs therein.
 - [3] A. Le Yaouanc, these proceedings and refs therein.
 - [4] B. Stech, Phys. Lett. B **354** (1995) 447.
 - [5] P. Ball and V. M. Braun, Phys. Rev. D **55** (1997) 5561.
 - [6] T. M. Aliev *et al.*, Phys. Rev. D **56** (1997) 4260.
 - [7] D. Becirevic, e-print [hep-ph/9707271](#).
 - [8] D. Melikhov, Phys. Rev. D **53** (1996) 2460; Phys. Lett. B **380** (1996) 363; e-print [hep-ph/9706417](#).
 - [9] N. Isgur and M. B. Wise, Phys. Lett. B **232** (1989) 113; *ibid.*, **237** (1990) 527; M. Luke, Phys. Lett. B **252** (1990) 447.
 - [10] N. Isgur and M. B. Wise, Phys. Rev. D **42** (1990) 2388.
 - [11] D. Melikhov, Phys. Lett. B **394** (1997) 385; D. Melikhov, N. Nikitin, S. Simula, e-print [hep-ph/9704268](#).
 - [12] D. Scora and N. Isgur, Phys. Rev. D **52** (1995) 2783.